

RESEARCH PAPER

Investigation of Some Autosomal Recessive Inherited Diseases (BLAD, DUMPS, CVM, and FXID) in Holstein Cattle

Döne TÜRKMEN¹ , Eymen DEMİR¹ , Ümit BİLGİNER¹ , Taki KARSLI^{2,*} 

¹Akdeniz University, Faculty of Agriculture, Department of Animal Science, Antalya, Republic of Türkiye

²Eskişehir Osmangazi University, Faculty of Agriculture, Department of Animal Science, Eskişehir, Republic of Türkiye

*Corresponding author

Article History

Received: .05 June.2023

Accepted: 20 December.2023

Corresponding Author*

E-mail: takikarsli@ogu.edu.tr

Keywords

Causative mutations

Genetic disorders

Genotyping

Molecular diagnosis

Malformation

Abstract

Genetic disorders are often transmitted via autosomal recessive inheritance, which negatively affect health, welfare, and yield traits in farm animals such as cattle. In this study, a total of 80 animals belonging to Holstein Friesian (HF) reared in the dairy farm of Akdeniz University were investigated in terms of Bovine Leukocyte Adhesion Deficiency (BLAD), Deficiency of Uridine Monophosphate Synthase (DUMPS), Factor XI Deficiency (FXID), and Complex Vertebral Malformation (CVM) via three molecular genotyping methods such as Polymerase Chain Reaction (PCR), PCR-Restriction Fragment Length Polymorphism (PCR-RFLP), and Allele-Specific PCR (AS-PCR). BLAD and DUMPS were analysed by PCR-RFLP, while PCR and AS-PCR were utilized to investigate FXID and CVM disorders, respectively. Based on fragment patterns on agarose gel electrophoresis, animals were genotyped as normal, mutant, and carrier. In this study, no carrier or mutant animals were detected for BLAD, CVM, DUMPS, and FXID diseases in HF cattle, since these animals were previously imported from disease-free dairy farms located in different provinces of Türkiye. Nevertheless, this population should be periodically checked for autosomal genetic disorders, since inherited diseases may be observed in the next generations due to causative mutations in the related genomic regions.

Introduction

Being able to survive in almost all climatic zones of the world, cattle may convert some crops and industrial products, which cannot be directly utilized for human nutrition, into animal-derived protein resources. As reported by the Food and Agriculture Organisation of the United Nations (FAO), 84% and 30% of the world's milk and beef production are met exclusively by cattle (FAO 2020). In Türkiye, cattle play a more important role in agriculture, since according to Turkish Statistical Institute (TUIK), a higher part of the milk (92.1%) and beef (74.8%) production are obtained from cattle farming (TUIK 2022a; 2022b). Hence, cattle are of great importance for human nutrition and the economies of developing countries such as Türkiye. It is known that a large part of the cattle population are exotic breeds (59.4%) and their crosses (42.8%), whereas 7.8% of the total population is predicted to be represented by native Turkish cattle breeds. Among exotic cattle populations, Holstein Friesian (HF) is the most reared breed in Türkiye (Demir et al., 2023).

It is essential to increase the quantity and quality of economically important traits via selection, while

health problems significantly decrease yield traits in livestock species. Therefore, the selection of healthy and tolerant animals is required to maintain the economic production systems. Inherited diseases negatively affect health and yield traits causing physical and functional anomalies in farm animals (Citek et al., 2006; 2007). Besides, as reported by Demir et al., (2021), genetic disorders are of negative effects on animal welfare.

Genetic disorders are mainly transmitted to the next generations via autosomal recessive inheritance in cattle (Windsor and Agerholm 2009). Using bulls with a high breeding value in artificial insemination increases the spreading risks of known and unknown genetic diseases in cattle breeding (Gentile and Testoni 2006; Citek and Blahova 2004). One of the most effective approaches to eliminating genetic diseases in a certain cattle population is to detect disease-carrier animals. However, due to the nature of the autosomal recessive inheritance, heterozygous animals, responsible for the occurrence of the diseases, show normal phenotype. This phenomenon requires accurate methods to detect

disease-carrier heterozygous animals. Thanks to molecular diagnosis methods, disease-carrier heterozygous animals could be detected even before birth in terms of numerous genetic disorders (Citek et al., 2006; 2007).

Bovine Leukocyte Adhesion Deficiency (BLAD), Deficiency of Uridine Monophosphate Synthase (DUMPS), Complex Vertebral Malformation (CVM), and Factor XI Deficiency (FXID) are frequently observed genetic disorders in HF breed (Meydan et al., 2010). BLAD disease is caused by a point mutation (A→G) (Shuster et al., 1992), at position 383 of the gene encoding the CD18 glycoprotein on the bovine chromosome 1 (Nagahata et al., 2004) and DUMPS disease is caused by another point mutation (C→T transition at codon 405) occurring in the UMPS gene on the same chromosome (Avanus ve Altinel 2017). The molecular basis of CVM disease is the result of a point mutation in nucleotide position 559 of the SLC35A3 gene located on bovine chromosome 3, which causes guanine to turn into thymine (Eren et al., 2019). FXID is caused by a 76 bp insertion in exon 12 of the Factor XI gene (Ghanem et al., 2005). Polymerase Chain Reaction - Restriction Fragment Length Polymorphism (PCR-RFLP) may detect BLAD and DUMPS diseases, while FXID and CVM could be assessed via PCR and Allele -Specific PCR (AS-PCR), respectively (Karslı et al 2011; Şahin et al., 2013). Hence, this study aims to screen 80 animals belonging to HF breed reared in the dairy farm of Akdeniz University in terms of some genetic disorders.

Materials and Methods

Ethics Statement

This research was approved by the Akdeniz University Animal Experiments Ethics Committee, Antalya, Türkiye (Protocol No: 1393/2022.01.005)

Sample Collection and DNA Extraction

In 2022, a total of 80 animals belonging to HF breed were randomly chosen from the dairy farm of the Akdeniz University, Faculty of Agriculture located in

Antalya province of Türkiye. Blood samples were taken from the jugular vein into vacuum tubes with K3EDTA as a coagulant. Blood samples were stored at -20°C until DNA extraction was performed. DNA was extracted via a salting-out method reported by Miller et al., (1988) following a spectrophotometer and 1% agarose gel electrophoresis in order to assess concentration and purity parameters.

Detection of BLAD, DUMPS, CVM, and FXID Genotypes

A total of three molecular methods such as PCR-RFLP (BLAD and DUMPS), AS-PCR (CVM), and traditional PCR (FXID) were utilized to genotype animals in terms of four genetic disorders. The methodology including oligonucleotide sequences as well as expected band sizes for normal, mutant, and carrier animals in terms of four genetic disorders was summarised in Table 1. Amplification was done in PCR reaction including 50 ng template DNA, 1.2 µL HQ buffer-GeneAll, 2 µL 10X buffer-GeneAll, 2.5 mM dNTPs, 10 pM primer, 2.5 U Taq DNA Polymerase, and 11.4 µL H₂O with the following protocol: 35 cycles of denaturation at 95 °C for 45 seconds, annealing at 52-62 °C for 45 seconds, and extension at 72 °C for 45 seconds, with an initial denaturation step at 95 °C for 5 minutes and a final extension step at 72 °C for 10 minutes. The annealing temperature was optimized at 57 °C, 52 °C, 55 °C, and 62 °C for BLAD, DUMPS, FXID, and CVM diseases, respectively. Since only one nucleotide differs in the primer sequence of the AS-PCR process, the annealing temperature was kept as high as possible to eliminate non-specific amplifications.

Results and Discussion

A 343 bp length PCR product cut by *TaqI* enzyme revealed two bands of 152 and 191 bp across all genotyped animals (Figure 1). No mutant or carrier was detected in the studied population in terms of BLAD disorder.

A total of three bands (53, 36, and 19 bp length) were observed in all animals in terms of DUMPS disease




Figure 1. Image of a 2% agarose gel showing the band lengths obtained after cutting PCR products for BLAD disease via the *TaqI* restriction enzyme. (DNA Ladder: 100 bp, Thermo, Catalog number: 15628050)



Figure 2. 3% agarose gel image of the band sizes obtained as a result of PCR-RFLP with *AvaI* restriction enzyme of PCR products for DUMPS disease (DNA Ladder: 100 bp, Thermo, Catalog number: 10416014)

(Figure 2). Due to failure in the discrimination power of agarose gel electrophoresis, the band of 19 bp is not clearly observable by bare eyes in Figure 2.

In terms of CVM disorder, PCR products were successfully amplified in all animals using normal primers, whereas no amplification was observed by using the mutant primer (Figure 3). This indicates that no CVM disease carrier is present in the studied cattle population.

All animals yielded a 244 bp length band in terms of FXID disorder, while the 230 bp length band was absent across the population. This finding indicates that there is no carrier or mutant animal in the studied cattle population.

Numerous studies showed that heterozygous carriers of the BLAD, DUMPS, CVM, and FXID disorders were present in HF breeds raised in different countries such as China (Zhang et al., 2010; Wang et al., 2011; Sun et al., 2011), Mexico (Virgen-Méndez et al., 2019), India (Patel et al., 2014), Poland (Gozdek et al., 2020). Similarly, Meydan et al., (2010) investigated the BLAD, DUMPS, CVM, Bovine Citrulinaemia (BC), and FXID diseases in 350 HF cattle reared in different regions of Türkiye in which a total of 14 and 12 animals were reported to be carriers of CVM and FXID disorders, respectively. On the contrary, no CVM carriers were reported in 150 heads of HF population raised in Kayseri province of Türkiye (Kulaklı and Akyüz 2011). Karsli et al.,

(2011) reported a low prevalence of FXID (0.4%), while no DUMPS carriers were present in a total of 504 animals belonging to HF breed reared in different districts of Antalya province of Türkiye. However, BLAD carriers were detected at the frequency of 2% and 4.8% in HF populations reared in Antalya and Kayseri provinces, respectively (Şahin et al., 2013; Akyüz et al., 2015). DUMPS and BC diseases were not reported in 219 Holstein cattle bred in Eskişehir, whereas two FXID and three BLAD carriers were present (Kaya et al., 2016). By using AS-PCR, Eren et al., (2019) confirmed that 7 out of 200 individuals analyzed were CVM carriers in HF population reared in Antalya. Screening a total of 48 animals in terms of BLAD and FXID disorders, Aksel et al., (2021) detected only one carrier animal for BLAD disorder.

Previous studies conducted on HF populations reared in Antalya province showed that the frequency of FXID carriers was 0.4%, while it was 2% for BLAD carriers, and 3.5% for CVM carriers (Karslı et al., 2011; Şahin et al., 2013). The frequency of FXID carriers in Holstein cattle reared in Burdur province was reported as 1.8% and the frequency of BLAD carriers as 2% (Ağaoğlu et al., 2015). Unlike previous researches, no carrier animals in terms of BLAD, FXID, and CVM disorders were detected in this study. It is thought that the main reason underlying this difference is the routine screening of artificial insemination bulls for these




Figure 3. Image of band sizes obtained in AS-PCR for CVM disease on a 2% agarose gel. (DNA Ladder: 100 bp, Thermo, Catalog number: 15628050)

Table 1. The method used to identify the relevant mutations and some descriptive information about genotyping

Diseases		BLAD	DUMPS	CVM	FXID
Method		PCR-RFLP		AS-PCR	PCR
Primer Sequence	Forward	CCTGCATCATATCC ACAG	GCAAATGGCTGAAGAAC ATTCTG	CACAATTTGTAGG TTCATGGCG	CCCACTGGCTAGGAA TCGTT
	Reverse	GTTTCAGGGGAAGA TGGAG	GCTTCTAACTGAACTCCT CGAGT	GTTATACTACAGG AGTCACCTCT	CAAGGCAATGTCATA TCCAC
	Mutant	-	-	CACAATTTGTAGG TTCATGCAT	-
PCR lengths		343	108	395	244 or 320
Restriction enzymes		<i>TaqI</i>	<i>AvaI</i>	-	-
Genotypes	Mutant	343	89 and 19	395	320
	Carrier	191, 152, and 343	89, 53, 36, and 19	395	244 and 320
	Normal	152 and 191	53, 36, and 19	395	244
References		Şahin et al., (2013)	Karşlı et al., (2011)	Eren et al., (2019)	Karşlı et al., (2011)

diseases for a long time, from which semen is obtained both abroad and locally. Another reason may be considered to be that the sampling was made from a single enterprise. However, as reported by Demir and Balcioglu (2019), HF population of the dairy farm of the Akdeniz University was re-established in 2017 by importing animals from several disease-free commercial enterprises located in ten different provinces of Türkiye. Therefore, the current population is thought to be a representative herd.

Conclusion

Genetic disorders in cattle, which are mostly transmitted through autosomal recessive inheritance, have negative impacts on animal health, welfare, and productivity. Molecular diagnosis methods are effective in detecting these disorders in carrier heterozygous animals, even before birth. In this study, no carrier animals in terms of BLAD, DUMPS, CVM, and FXID disorder were detected in HF population raised on the farm of the Akdeniz University due to the fact that these animals were imported from disease-free farms localized in ten provinces in Türkiye. This information can be used to improve breeding strategies, including the selection of animals with high breeding values and the prevention of the spreading of known and unknown genetic diseases in cattle breeding. Further studies are recommended to extend the screening to other cattle populations in different regions of Turkey with larger sample sizes to understand the distribution of these genetic disorders and develop strategies for the prevention and control of these diseases.

Acknowledgments

This work was supported by the Scientific and Technological Research Council of Türkiye (Project No: 1919B012101334), Republic of Türkiye.

Conflict of Interest

The authors declare no conflicts of interest.

References

- Ağaoğlu, Ö. K., Ağaoğlu, A. R., & Saatci, M. (2015). Estimating allele frequencies of some hereditary diseases in Holstein cattlereared in Burdur Province, Türkiye. *Turkish Journal of Veterinary & Animal Sciences*, 39(3), 338-342. doi: 10.3906/vet-1412-13
- Akyuz, B., Sariozkan, S., & Bayram, D. (2015). The prevalence of BLAD and comparison of some production parameters in carrier and non-carrier Holstein cows in Kayseri province, Türkiye. *Journal of Faculty of Veterinary Medicine, Erciyes University/Erciyes Üniversitesi Veteriner Fakültesi Dergisi*, 12(1), 31-35.
- Avanus, K., & Altinel, A. (2017). Inherited diseases of Holstein cattle: Story so far in Turkey. *Journal of Istanbul Veterinary Sciences*, 1(2), 40-46. doi:10.30704/http-www-jivs-net.324403
- Citek, J., & Blahova, B. (2004). Recessive disorders-a serious health hazard?. *Journal of Applied Biomedicine*, 2, 187-194. doi:10.32725/jab.2004.022
- Citek, J., Rehout, V., Hajkova, J., & Pavkova, J. (2006). Monitoring of the genetic health of cattle in the Czech Republic. *Veterinární Medicína Czech*; 51(6): 333-339. doi: 10.17221/5553-vetmed
- Citek, J., Rehout, V., Vecerek, L., & Hajkova, J. (2007). Genotyping glycogen storage disease type II and type V in cattle reared in the Czech Republic. *Journal of Veterinary Medicine*, 54, 257-259. doi: 10.1111/j.1439-0442.2007.00931.x
- Demir, E., & Balcioglu, M. S. (2019). Genetic diversity and population structure of four cattle breeds raised in Turkey using microsatellite markers. *Czech Journal of Animal Science*, 64(10), 411-419. doi: 10.17221/62/2019-cjas
- Demir, E., Bilginer, U., Balcioglu, M. S., & Karşlı, T. (2021). Direct and indirect contributions of molecular genetics to farm animal welfare: a review. *Animal Health Research Reviews*, 22(2), 177-186. doi: 10.1017/S1466252321000104
- Demir, E., Moravčíková, N., Kaya, S., Kasarda, R., Doğru, H., Bilginer, Ü., Balcioglu, M.S., & Karşlı, T. (2023). Genome-wide genetic variation and population structure of native and cosmopolitan cattle breeds reared in Türkiye. *Animal Biotechnology*, 1-10. doi: 10.1080/10495398.2023.2235600

- Eren, M. G., Balcioglu, M. S., & Demir, E. (2019). Detection of complex vertebral malformation in Holstein cattle reared in Antalya using allele specific PCR. *Mediterranean Agricultural Sciences* 32(3): 443-446 doi:10.29136/mediterranean.578015
- FAO, FAOSTAT, (2022). Food and Agriculture data, Crops and livestock products, <https://www.fao.org/faostat/en/#data/QCL>, Date of access: 03.06. 2022
- Gentile, A., & Testoni, S. (2006). Inherited disorders of cattle. *Slovenian Veterinary Research*, 43(1), 17-29.
- Gozdek, M., Kolenda, M., Kamola, D., & Sitkowska, B. (2020). Report on the incidence of hereditary disorders (BLAD, DUMPS) in the Polish population of Holstein-Friesian cattle, *Acta Scientiarum Polonorum. Zootechnica*, 19(3),15-22. doi:10.21005/asp.2020.19.3.02
- Karsli, T., Sahin, E., Karsli, B.A., Alkan, S., Balcioglu, M.S. (2011). Identification of alleles for factor XI (FXID) and uridine monophosphate synthase (DUMPS) deficiencies in Holstein cows reared in Antalya. *Kafkas Universitesi Veteriner Fakültesi Dergisi* 17, 503–505.
- Kaya, M., Meydan, H., Kıyma, Z., Alan, M., & Yildiz, M. A. (2016). Screening for bovine leukocyte adhesion deficiency, deficiency of uridine monophosphate synthase, bovine citrullinaemia and factor XI deficiency in Holstein cattle. *Indian Journal of Animal Sciences*, 86(8), 900-903.
- Kulaklı, G. N., & Akyüz, B. (2011). Kayseri bölgesinde yetiştirilen Holştayn sığırlarında kompleks vertebral malformasyon hastalığı geninin allel frekansının belirlenmesi. *Erciyes Üniversitesi Veteriner Fakültesi Dergisi*, 8(2), 69-74.
- Kunieda, T. (2005). Identification of genes responsible for hereditary diseases in Japanese beef cattle. *Animal Science Journal*, 76, 525–533. <https://doi.org/10.1111/j.1740-0929.2005.00300.x>
- Meydan, H., Yildiz, M. A., & Agerholm, J. S. (2010). Screening for bovine leukocyte adhesion deficiency, deficiency of uridine monophosphate synthase, complex vertebral malformation, bovine citrullinaemia, and factor XI deficiency in Holstein cows reared in Türkiye. *Acta Veterinaria Scandinavica*, 52, 1-8. doi: 10.1186/1751-0147-52-56.
- Miller, S., Dykes, D., & Plesky, H.A. (1988). Simple salting out procedure for extracting DNA from human cells. *Nucleic Acids Research*, 16: 1215. doi: 10.1093/nar/16.3.1215.
- Nagahata, H. (2004). Bovine leukocyte adhesion deficiency (BLAD): a review. *Journal of Veterinary Medical Science*, 66(12), 1475-1482. doi: 10.1292/jvms.66.1475
- Patel, R.K., & Patel, A.K. (2014). Comparative review of recessive genetic disorders occurrence in Indian Cattle. *Current Trends in Biotechnology and Pharmacy*, 8(3), 321-335.
- Şahin, E., Karsli, T., Galiç, A., & Balcioglu, M.S. (2013). Identification of bovine leukocyte adhesion deficiency (BLAD) and bovine citrullinaemia (BC) alleles in Holstein cows reared in Antalya region. *Journal of Applied Animal Research*, 41(1): 56-60. doi:10.1080/09712119.2012.738221
- Schwenger, B., Tammen, I., Aurich, C. (1993). Detection of homozygous recessive genotype for deficiency of uridine monophosphate synthase by DNA typing among bovine embryos produced in vitro. *Journal of Reproduction & Infertility*, 100, 511–514. doi: 10.1530/jrf.0.1000511.
- Shuster, D.E, Kehrlı, M., Ackermann, M.R., & Gilbert, R. (1992). Identification and prevalence of a genetic defect that causes leukocyte adhesion deficiency in Holstein cattle. *Proceedings of the National Academy of Sciences*, 89, 9225-9229. doi: 10.1073/pnas.89.19.9225
- Sun, D. X., Fan, X.H., Xie, Y., Chu, Q., Sun, Y., Zhang, Y., & Shi, W. H. (2011). Distribution of recessive genetic defect carriers in Chinese Holstein. *Journal of dairy science*, 94(11), 5695-5698. doi: 10.3168/jds.2011-4345.
- TUIK, 2022a, Türkiye İstatistik Kurumu, Çiğ Süt Üretim İstatistikleri 2020-2021, <https://data.tuik.gov.tr/Bulten/Index?p=Cig-Sut-Uretim-Istatistikleri-2020-2021-45861>, Date of Access: 03.06. 2022.
- TUIK, 2022b, Türkiye İstatistik Kurumu, İstatistik Tablolar, Hayvansal Üretim İstatistikleri (Yıllık), Büyükbaş Hayvan Sayıları, <https://data.tuik.gov.tr/Bulten/Index?p=Hayvansal-Uretim-Istatistikleri-Aralik-2021-45593>, Date of Access: 03.06. 2022.
- TUIK, 2022c, Türkiye İstatistik Kurumu, Kırmızı Et Üretim İstatistikleri 2020-2021, <https://data.tuik.gov.tr/Bulten/Index?p=Kirmizi-Et-Uretim-Istatistikleri-2020-2021-45671>, Date of Access: 03.06. 2022.
- Virgen-Méndez, A., Ayala-Valdovinos, M.A., Galindo-García, J., Sánchez-Chiprés, D.R., Lemus-Flores, C., & Duifhuis-Rivera, T. (2019). Carrier frequency of autosomal recessive disorders (BC, BLAD, FXID and CVM) in Holstein cows in Jalisco, Mexico. *Pesquisa Veterinária Brasileira*, 39, 481-484. doi: 10.1590/1678-5150-PVB-6202
- Wang, C., Tong, Q., Hu, X. Z., Yang, L. G., Zhong, X. Q., Yu, Y., Wu, J.J., Liu, W.J., Li, X., Hua, G.H., Zhao, H.Q., & Zhang, S. J. (2011). Identification of complex vertebral malformation carriers in Holstein cattle in south China. *Genetics and Molecular Research*, 10(4), 2443-2448. doi:10.4238/2011.October.13.1.
- Windsor, P.A., & Agerholm, J.S. (2009). Inherited diseases of Australian Holstein-Friesian cattle. *Australian Veterinary Journal*, 87, 193–199. doi: 10.1111/j.1751-0813.2009.00422.x.
- Zhang, K., Wang, Z., & Wang, Q. (2010). Detecting factor XI deficiency in Holstein cattle using PCR analysis. *Agricultural Science & Technology-Hunan*, 11(5), 109-111.