



RESEARCH ARTICLE

The Effects of Different Doses of eCG Injection on Reproductive Parameters in Nulliparous Merino Ewes During the Non-Breeding Season

Neffel Kürşat AKBULUT^{1,*}  Metehan KUTLU¹ 

¹Department of Obstetrics and Gynaecology, Faculty of Veterinary Medicine, Necmettin Erbakan University, Ereğli, Konya, Türkiye.

Article History

Received: 25 Sep 2024

Accepted: 30 Oct 2024

Corresponding Author

E-mail: nkakbulut@gmail.com

Keywords

Conception rate

eCG

Medroxyprogesterone acetate

Pregnancy rate

Abstract

In this study, the effects of adding 400, 500, and 600 IU eCG to progesterone treatment on rate of estrus, conception, pregnancy, fetal mortality, lambing, multiple birth and litter size in nulliparous Merino ewes the out of breeding season were investigated. A total of 90 nulliparous Merino ewes were utilized in the study. On day 0, a vaginal sponge containing 60 mg medroxyprogesterone acetate was inserted for a seven-day period. Ewes were randomly divided into three groups at the time of sponge removal. Group I (G-400), comprising 30 ewes, received 400 IU of eCG. Group II (G-500) received 500 IU of eCG (n=30), while Group III (G-600) received 600 IU of eCG (n=30). The statistical analysis revealed that the conception rates between the groups were statistically higher in the G-600 group than in the G-400 and G-500 groups (G-400 vs. G-600, P=0.091; G-500 vs. G-600, P=0.080). Pregnancy rates between the groups were found to be statistically higher in the G-600 group than in the G-400 group (P<0.021). The administration of a high dose of eCG (600 IU) in conjunction with progesterone synchronization yielded favorable outcomes with respect to reproductive parameters, particularly an elevated conception and pregnancy rates.

Introduction

In sheep breeding, the primary objectives are to achieve greater efficiency without increasing costs and to enhance reproductive performance (Özyurtlu and Bademkiran, 2010). The profitability of sheep farming is closely linked to the number of lambs produced annually (Bulbul *et al.*, 2014). Exogenous hormones are the leading biotechnological products used in animal husbandry, especially to improve fertility. During the breeding season, progesterone and and prostaglandin F₂ alpha (PGF₂α) can be used for estrus synchronization and equine chorionic gonadotropin (eCG) to induce ovulation in sheep farming (Köse *et al.*, 2022). The protocols entail the administration of progesterone to the ewe, whether short- or long-term, with the objective of stimulating the luteal phase of the cycle. Once this period has elapsed, the source of the

progesterone is removed and eCG is applied (Cline *et al.*, 2001).

eCG is a glycoprotein hormone secreted by trophoblastic epithelial cells of fetal origin that form the equine endometrial cups (Manteca Vilanova *et al.*, 2019). eCG has been observed to exhibit high affinity for follicle-stimulating hormone (FSH) and luteinizing hormone (LH)-related receptors. This affinity has been associated with increased estradiol before ovulation (Rostami *et al.*, 2011) and increased progesterone after ovulation in dairy cows (Rowe *et al.*, 2019). In these treatments, the estrus response and onset of estrus may be affected by factors such as the dose and the time of administration of eCG, breed and season, body condition and geographical region (Quintero-Elisea *et al.*, 2022).

In estrous synchronization protocols, low doses of eCG generally reduce the time to onset of estrus.

Additionally, evidence indicates that low doses of eCG typically enhance the number and size of large follicles, though it may exert minimal influence on the ovulation rate (Moakhar *et al.*, 2012). The effect of low-dose eCG in non-breeding season ewes is of great interest. The effect of eCG on the dynamics of the follicular wave in ewes during the non-breeding season is well known (Roshan *et al.*, 2023). A deeper understanding of this mechanism could facilitate the development of more effective methods for regulating sheep breeding (Barrett *et al.*, 2004).

The number of progesterone and eCG-based studies on nulliparous ewes is insufficient, and the available information is limited (Santos-Jimenez *et al.*, 2022). In studies conducted out of the breeding season, there are reports indicating that multiparous ewes have show a better response to these hormones compared to nulliparous ewes (Santos-Jimenez *et al.*, 2022), as well as studies reporting no significant difference (Ungerfeld and Rubianes, 1999). In a study in ewe lambs, higher pregnancy and conception rates were found in the groups using eCG compared to the control group (Dias *et al.*, 2020). It can be posited that the utilisation of eCG in ewe lambs may facilitate the attainment of a greater number of lambs.

It is thought that increasing the dose of eCG may induce estrus by stimulating follicular development and improve some reproductive parameters. It is hypothesized that the response to progesterone and different doses of eCG may vary non-breeding season in nulliparous Merino ewes. In this study, the effects of adding 400, 500, and 600 IU eCG to progesterone treatment on rate of estrus, conception, pregnancy, fetal mortality, lambing, multiple birth and litter size in nulliparous Merino ewes out of the breeding season were investigated.

Materials and Methods

The present study was conducted with approval from Selçuk University Animal Experiments Local Ethics Committee, Konya, Türkiye (2024/084).

Animals

This study was conducted in a commercial sheep farm (Lat: 37° 86' 44.06» N, Long: 34° 16' 33.55» E and Alt: 1.020 m) in Konya province, Türkiye during the non-breeding season in 2024. A total of 96 nulliparous (1 year old, 45-50 kg) Merino ewes, clinically healthy with no abnormalities in the perineum, were used. Six ewes whose sponge had been fell were excluded from the study. The study continued with 90 ewes. The ewes were permitted to graze on pasture for a duration of 12 hours each day and were not provided with any compound feed. They had ad libitum access to water. Prior to mating, the animals were not subjected to any nutritional flushing.

Treatment groups, synchronization and mating protocols

On day 0, a vaginal sponge containing 60 mg of medroxyprogesterone acetate (Esponjavet®, Hipra, Spain) was inserted for a seven-day period. On day 7, the sponges were removed. The ewes were randomly divided into 3 groups at the time the sponges were removed and 400 IU of eCG (Oviser®, Hipra, Spain) was injected to ewes in Group 1 (G-400) (n=30), 500 IU of eCG to ewes in Group 2 (G-500) (n=30), and 600 IU of eCG was injected to the ewes in Group 3 (G-600) (n=30).

A teaser ram was employed twice daily for 1-hour sessions over 36 hours following the removal of sponges to detect estrus. Ewes identified as being in estrus were hand-mated with one of the proven fertile Merino rams, maintaining a ewe-to-ram ratio of 7:1.

Ultrasonography and calculation of reproductive parameters

In all ewes, transabdominal ultrasound examination (Hitachi EUB-405, Japan, 3.5 MHz convex probe) was performed to diagnose pregnancy on day 45 post-mating. Pregnancy (single/twin) were diagnosed when embryos were seen on ultrasound examination. The litter size was determined at parturition.

The rate of estrus, conception, pregnancy, fetal mortality, lambing, multiple birth and litter size were calculated as reproductive parameters as follows;

$$\text{Estrus rate} = \frac{\text{the number of ewes showing estrus behaviours}}{\text{the number of ewes receiving intravaginal sponge}}$$

$$\text{Conception rate} = \frac{\text{the number of pregnant ewes}}{\text{the number of mated ewes}} \times 100$$

$$\text{Pregnancy rate} = \frac{\text{the number of pregnant ewes}}{\text{the number of ewes receiving intravaginal sponge}}$$

$$\text{Fetal mortality rate} = \frac{\text{number of detected death fetus}}{\text{the number of pregnant ewes}} \times 100$$

$$\text{Lambing rate} = \frac{\text{the number of lambing ewes}}{\text{the number of pregnant ewes}} \times 100$$

$$\text{Multiple birth rate} = \frac{\text{the number of multiple lambing ewes}}{\text{the number of pregnant ewes in each group}} \times 100$$

$$\text{Litter size} = \frac{\text{the number of total lambs}}{\text{the number of lambing ewes}} \times 100$$

Statistical analysis

Statistical analyses were performed using SAS (Version 8.0). Reproductive parameters were assessed using the Chi-squared test, Fisher's exact test (Selvi, 2024), and PROC GENMOD procedure. The results were reported as percentages. Statistical significance was defined as $p < 0.05$, and tendency was considered for $0.05 < p \leq 0.10$.

Results

Results for the rate of estrus, conception, pregnancy, fetal mortality, lambing, multiple birth and litter size are given in Table 1.

In the statistical analysis, differences between the groups in rate of estrus, fetal mortality, lambing, multiple birth and litter size were found to be insignificant ($P > 0.05$). Conception rates between groups were found to be higher in the G-600 group than in the G-400 and G-500 groups (G-400 vs G-600 $P=0.091$; G-500 vs G-600 $P=0.080$). Pregnancy rates between the groups were found to be statistically higher in the G-600 group than in the G-400 group ($P<0.05$).

Discussion

One method for increasing productivity in sheep is to obtain offspring during the anestrous period in Türkiye. During the non-breeding season, progestagens and eCG hormones are often used to stimulate functional activity in the ovaries. (Kaçar *et al.*, 2008). eCG is known to be the most commonly used hormone to improve fertility in reproduction (Moakhar *et al.*, 2012). The effectiveness of eCG is affected by many factors such as injection method, dose and source of eCG and season (Ali, 2007).

The objective of the presented study was to investigate the effects of different doses of eCG on reproductive parameters in nulliparous Merino ewes that had been synchronized with progesterone during the non-breeding season. In ewes, it has been reported that short term intravaginal progestagen treatment successfully induce and synchronize estrus in both

breeding and non-breeding seasons (Ahmed Amer and Maher Hazzaa, 2009; Ataman *et al.*, 2006). A study conducted non-breeding season demonstrated that the administration of eCG in conjunction with short-term progesterone resulted in a high estrus rate in both nulliparous and multiparous ewes (Ungerfeld and Rubianes, 1999). Santos-Jiménez (2022) observed that the estrus rate was higher in multiparous ewes than in nulliparous ewes during the non-breeding season. This result was attributed to the silent estrus of nulliparous ewes.

In some studies, conducted during the non-breeding season, estrus rates were observed to be high in ewes that had been synchronized with progesterone-based eCG applications at varying doses (Efe, 2010; Ungerfeld and Rubianes, 1999). The administration of 300, 500, and 700 IU of eCG following the intravaginal sponge application for 14 days during the seasonal anoestrus period resulted in estrus rates of 69.66% in all groups of Algerian Rembi sheep. (Bacha *et al.*, 2014). Aköz *et al.* (2006) injected 300, 500 and 700 IU eCG after using intravaginal sponge containing 30 mg and 40 mg fluorogesterone acetate (FGA) in Merino x Akkaraman crossbred ewes during non-breeding season, and found the estrus rates to be 100%, 93.3% and 100% in the 30 mg FGA group, and 93.3%, 92.8% and 100% in the 40 mg FGA group, respectively. As with the aforementioned studies, it can be stated that the administration of varying doses of eCG does not result in statistically significant alterations in estrus rates, as observed in the presented study. It has been reported that eCG exhibits both FSH and LH activity and that these hormones are necessary for peri-ovulatory maturation of mammalian follicles (Murphy, 2018). Additionally, some studies have reported that eCG

Table 1. Reproductive parameters of nulliparous ewes.

	G-400 (n=30)	G-500 (n=30)	G-600 (n=30)	P
Estrus rate (%)	73.3 (22/30)	90.0 (27/30)	90.0 (27/30)	0.135
Conception rate (%)	59.1 ^c (13/22)	53.3 ^c (16/27)	81.5 ^d (22/27)	**
Pregnancy rate (%)	43.3 ^a (13/30)	53.3 ^{ab} (16/30)	73.3 ^b (22/30)	*
Fetal mortality rate (%)	7.6 (1/13)	18.8 (3/16)	13.6 (3/22)	0.679
Lambing rate (%)	92.3 (12/13)	81.2 (13/16)	86.3 (19/22)	0.679
Multiple birth rate (%)	16.7 (2/12)	23.1 (3/13)	42.1 (8/19)	0.261
Number of Lambs	14	16	28	
Single	10	10	11	
Twin	2 (4)	3 (6)	7 (14)	
Triplets	-	-	1 (3)	
Litter Size	1.17 (14/12)	1.23 (16/13)	1.47 (28/19)	0.244

Notes: Values with different superscripts (a, b) in the same row are significant differences among the groups ($P < 0.05$)

Values with different superscripts (c, d) in the same row are tendency differences ($0.05 < P \leq 0.1$) among the groups.

* Pregnancy rate P value G-400 vs G-600 $P=0.021$

**Conception rate P value G-400 vs G-600 $P=0.091$; G-500 vs G-600 $P=0.080$

The numbers in parentheses are the formulation of the reproductive parameter.

causes an increase in progesterone concentrations (Baruselli *et al.*, 2009). The findings of this study indicate that the administration of 600 IU eCG may enhance luteal activity, thereby improving pregnancy rates. While numerical differences were observed in certain examined characteristics, no statistically significant differences were identified. Takcı *et al.* (2023) reported that different eCG doses did not have a significant effect on reproductive parameters in primiparous ewes during the early anoestrus period. Köse *et al.* (2022) reported that different doses (400 and 600 IU) of eCG did not result in any differences in reproductive performance in ewes during the breeding season. Efe (2010) stated that although the use of 500 - 600 and 700 IU eCG after progesterone during the anoestrus period in ewes caused higher pregnancy rates than the control group (without eCG) and the dose difference did not affect the pregnancy rates. In a study conducted on Merino ewes during the non-breeding season, Doğanay (2011) used 400 and 600 IU eCG in addition to 14-day intravaginal progesterone application and found no statistical difference in pregnancy rates. Koyuncu *et al.* (2001) stated that the use of 0, 500 and 700 IU eCG in ewes synchronized with progesterone and found the litter size to be 1.21, 1.58 and 1.96, respectively on Kivırcık ewes during the breeding season. In studies on the effects of eCG on reproductive parameters in ewes, it is seen that many factors are effective and therefore the results of the studies are not consistent (Köse *et al.*, 2016). In order to eliminate this confusion in pregnancy rates, the optimum eCG dose needs to be determined. (Sharif *et al.*, 2023). For this purpose, Sharif *et al.* (2023) used 300 and 600 IU eCG excluding the control group (without eCG) in Beetal goats at the beginning of the breeding season, and reported an increase in pregnancy rates in the group receiving 600 IU eCG compared to the other groups. Moaktar *et al.* (2012) in their study on Chall sheep during the natural breeding season, used 0, 450, 550, 650, 750, and 850 IU eCG and as a result, obtained higher pregnancy rates in the groups that received 550 and 650 IU eCG. Based on the studies conducted, it has been observed that different reproductive outcomes occur during breeding and non-breeding season. Many authors have argued that these results are due to the different responses of sheep breeds to eCG doses (Quintero-Elisea *et al.*, 2022; Roshan *et al.*, 2023). In our study, the Merino ewes used are capable of exhibiting reproductive activity for most of the year and have good fertility rates and, it is thought that the reproductive outcomes obtained, compared to other studies, to be due to the breed-specific characteristics.

Conclusion

In conclusion, it was determined that high dose eCG (600 IU) used in addition to synchronization with intravaginal progesterone during non-breeding season in nulliparous Merino ewes had positive results on

reproductive parameters, especially causing an increase in pregnancy and conception rates.

Conflict of Interest

The authors declare that there is no conflict of interest.

References

- Ahmed Amer, H., Maher Hazzaa, A. (2009). The effect of different progesterone protocols on the reproductive efficiency of ewes during the non-breeding season. *Veterinarski arhiv*, 79: 19-30.
- Akoz, M., Bulbul, B., Bozkurt A, M., Dere, S. (2006). Induction of multiple births in akkaraman cross-bred sheep synchronized with short duration and different doses of progesterone treatment combined with pmsg outside the breeding season. *Bulletin-Veterinary Institute in Pulawy*, 50: 97.
- Ali, A. (2007). Effect of time of eCG administration on follicular response and reproductive performance of fga-treated ossimi ewes. *Small Ruminant Research*, 72: 33-37.
- Ataman, M., Akoz, M., Akman, O. (2006). Induction of synchronized oestrus in akkaraman cross-bred ewes during breeding and anestrus seasons: The use of short-term and long-term progesterone treatments. *Revue de médecine vétérinaire*, 157: 257-260.
- Bacha, S., Khiati, B., Hammoudi, S. M., Kaidi, R., Ahmed, M. (2014). The effects of dose of pregnant mare serum gonadotropin (pmsg) on reproductive performance of algerian rembi ewes during seasonal anoestrus. *Journal of Veterinary Science and Technology*, 5: 190.
- Barrett, D., Bartlewski, P., Batista-Arteaga, M., Symington, A., Rawlings, N. (2004). Ultrasound and endocrine evaluation of the ovarian response to a single dose of 500 iu of eCG following a 12-day treatment with progestogen-releasing intravaginal sponges in the breeding and nonbreeding seasons in ewes. *Theriogenology*, 61: 311-327.
- Baruselli, P. S., Ferreira, R. M., Sá Filho, M. F., Nasser, L. F., Rodrigues, C. A., Bó, G. A. (2009). Bovine embryo transfer recipient synchronisation and management in tropical environments. *Reproduction, Fertility and Development*, 22: 67-74.
- Bulbul, B., Kirbas, M., Aktas, A. H., Kose, M., Ataman, M. B., Coyan, K., Kan, M. (2014). Investigation of accelerated lambing possibility of anatolian merino sheep.
- Cline, M., Ralston, J., Seals, R., Lewis, G. (2001). Intervals from norgestomet withdrawal and injection of equine chorionic gonadotropin or pg 600 to estrus and ovulation in ewes. *Journal of Animal Science*, 79: 589-594.
- Dias, J., Miranda, V., Oliveira, F., Junior, S. V., Haas, C., Costa, V., Lucia Jr, T., Vieira, A., Corcini, C., Gasperin, B. (2020). Treatment with eCG and hCG to induce onset of estrous cycles in ewes during the non-breeding season: Effects on follicular development and fertility. *Animal reproduction science*, 212: 106232.
- Doğanay, M. (2011). The impact of different dosage usage of progestagen+pmsg on the stimulation of ovarian

- activity, synchronication of sexual cycles and subsequent fertility rate of merino ewes during the anoestros season Master Thesis, Health Sciences Institute, Ankara University.
- Efe, A. (2010). Effects of different doses of pregnant mare serum gonadotrophine on reproduction in anoestrus sheep. Master Thesis, Yüzüncü Yıl University.
- Kaçar, C., Kamiloğlu, N., Pancarcı, Ş., Güngör, Ö., Güvenç, K., Gürbulak, K., Saban, E. (2008). The effect of administration of testosterone antibody, β -carotene and vitamin e on multiple pregnancy and mda (malondialdehyde) in tuj breed sheep in non-breeding season. Kafkas Üniversitesi Veteriner Fakültesi Dergisi, 14.
- Koyuncu, M., Uzun, Ş. K., Şengül, L. (2001). Synchronization of oestrus and the possibilities of improving reproductive performance by using progestagen and different doses of pmsg in kivircik ewes. Turkish Journal of Veterinary & Animal Science, 25: 971-974.
- Köse, M., Bayrıl, T., Küçükaslan, İ., Bademkiran, S., Özyurtlu, N., Atlı, M., Yıldız, A. Ş. (2022). Effect of different doses of equine chorionic gonadotropini on some fertility parameters in zom ewes synchronized with progesterone and prostaglandin f2 alfa during breeding season. Dicle University Journal of Faculty of Veterinary Medicine, 15: 99-103.
- Köse, M., Dursun, Ş., Bülbül, B., Kırbaş, M., Demirci, U. (2016). Investigation of possibility of increasing lamb production with flushing plus ram effect or the administration of various pregnant mare serum gonadotropin doses in akkaraman ewes. Veterinary Sciences and Practices, 11: 54-59.
- Manteca Vilanova, X., De Briyne, N., Beaver, B., Turner, P. V. (2019). Horse welfare during equine chorionic gonadotropin (ecg) production. Animals, 9: 1053.
- Moakhar, H. K., Kohram, H., Shahneh, A. Z., Saberifar, T. (2012). Ovarian response and pregnancy rate following different doses of ecg treatment in chall ewes. Small Ruminant Research, 102: 63-67.
- Murphy, B. (2018). Equine chorionic gonadotropin: An enigmatic but essential tool. Animal Reproduction (AR), 9: 223-230.
- Özyurtlu, N., Bademkiran, S. (2010). Estrus synchronization and induction of estrus methods in sheep. Dicle University Journal of Faculty of Veterinary Medicine, 1: 17-22.
- Quintero-Elisea, J. A., Olguín-Arredondo, H. A., Velázquez-Morales, J. V., Garay-Martínez, J. R., Lizeth, V.-R., Limas-Martínez Andrés, G., Joaquín-Cancino, S. (2022). Effect of breed, breeding season, ecg dose, and ecg application time on the estrous cycle of hair ewe lambs. Agro Productividad, 15.
- Roshan, N. J., Garoussi, M. T., Akbarinejad, V. (2023). Evaluation of the effect of melatonin implantation in rams and ecg dose in ewes synchronized by a cidr-ecg protocol on reproductive performance of lacaune sheep breed during non-breeding season. Animal reproduction science, 259: 107365.
- Rostami, B., Niasari-Naslaji, A., Vojgani, M., Nikjou, D., Amanlou, H., Gerami, A. (2011). Effect of ecg on early resumption of ovarian activity in postpartum dairy cows. Animal reproduction science, 128: 100-106.
- Rowe, S., Pryor, L., Tranter, W., Hosie, J., Cavalieri, J. (2019). Effect of equine chorionic gonadotropin on reproductive performance in a dairy herd in northern queensland, australia. Theriogenology, 125: 30-36.
- Santos-Jimenez, Z., Martínez-Ros, P., Encinas, T., Morales-Cruz, J. L., Guerrero-Gallegos, H. Z., Gonzalez-Avalos, R., Gonzalez-Bulnes, A., Guillen-Muñoz, J. M. (2022). Ovarian response and fertility after short-term progestagen/ecg treatments are compromised in nulliparous sheep during non-breeding season. Veterinary Sciences, 9: 663.
- Selvi, M. H. (2024). The use of statistics in veterinary sciences and the test methods used. Research and Practice in Veterinary and Animal Science, 1: 43-50.
- Sharif, B., Hassan, M., Arshad, U., Tahir, M. Z., Ahmad, E., Khan, M. I., Shahzad, M., Mohsin, I., Sosa, F., Rehman, A. (2023). Effect of ecg dose on ovarian haemodynamics, hormonal profiles and prolificacy rate when oestrus was induced during low-breeding season in beetal goats. Reproduction in Domestic Animals, 58: 48-59.
- Takcı, A., Kıvrak, M. B., Yüksel, M. (2023). Determination of the effect of different doses of ecg applications on reproductive parameters in primiparous akkaraman kangal sheep. Turkish Journal of Agriculture-Food Science and Technology, 11: 1128-1133.
- Ungerfeld, R., Rubianes, E. (1999). Effectiveness of short-term progestogen primings for the induction of fertile oestrus with ecg in ewes during late seasonal anoestrus. Animal Science, 68: 349-353.