

# The Occurrence of Arsenic, Cadmium, and Lead Residues in Cattle Feed Collected in Kırıkkale, Türkiye: A Preliminary Study

Erkan Şahin<sup>1</sup>, Hatice Akay Sarıkaya<sup>2</sup>, Ebru Yıldırım<sup>3</sup>, Miyase Çınar<sup>4,\*</sup>

<sup>1</sup> Kırıkkale University, Institute of Health Sciences, Biochemistry, Kırıkkale, Türkiye

<sup>2</sup> International Center for Livestock Research and Training, Türkiye

<sup>3</sup> Department of Pharmacology and Toxicology, Faculty of Veterinary Medicine, Kırıkkale University, Kırıkkale, Türkiye

<sup>4</sup> Department of Biochemistry, Faculty of Veterinary Medicine, Kırıkkale University, Kırıkkale, Türkiye

\*Corresponding Author

## Article History

Received: 28 Nov 2021

Accepted: 28 Sep 2022

First Online: 29 Oct 2022

## Corresponding Author\*

Tel.: 03183574242/6560

E-mail: miyasecinar@kku.edu.tr

## Keywords

Arsenic

Cadmium

Cattle feed

Heavy metal

Lead

## Abstract

The aim of this study is to investigate the occurrence of arsenic (As), cadmium (Cd), and lead (Pb) residues in 15 cattle feed collected from dairy and beef cattle breeding farms located in 6 different districts (Kırıkkale Province, Yahşihan, Karakeçili, Balışeyh, Delice and Keskin) of Kırıkkale. The quantitative analysis of the As, Cd and Pb levels of the samples was analyzed by ICP-MS. Arsenic and Pb residues were found in all of the collected samples, while Cd residues were found in 13 (86. 67%) of the collected samples. In the samples with heavy metal residues; the mean value  $\pm$  standard deviation for As, Cd, and Pb were found as  $0.1475 \pm 0.1060$  mg/kg,  $0.0382 \pm 0.0079$  mg/kg and  $0.1944 \pm 0.1074$  mg/kg respectively. The maximum tolerable limit in cattle feed is 2 mg/kg for As, 5 mg/kg for Pb, and for Cd in cattle feed is 1 mg/kg for Cd according to an official announcement (2014/11) of the Ministry of Agriculture and Forestry Turkey. The results of this study showed that detected As, Cd and Pb levels in cattle feeds were below the tolerable limit. Therefore, it is suggested that the heavy metal pollution in terms of As, Cd, and Pb in feed does not pose a risk for cattle farming in Kırıkkale province.

## Introduction

The term heavy metal can be used for elements having an atomic mass of 200 or more like mercury, thallium, lead and bismuth. In practice, heavy metals include metals that is clinically cause undesirable effects and potentially dangerous for health (Baldwin and Marshall, 1999). Some metals such as cobalt, copper, chromium, and iron are necessary to maintain many biochemical and physiological processes in living organisms including animals and plants. On the other hand, metals such as As, Cd, Pb, and mercury (Hg) do not play a role in biological functions, and these undesirable substances in animal feeds are called contaminants. These metals threaten public health at high doses, and may even cause organ damage at low doses (Hejna *et al.*, 2018). It is known that especially Cd, Pb and less

commonly Hg cause kidney damage (Garcia *et al.*, 2018). There are also findings reporting that these heavy metals cause cardiovascular diseases and chronic heart diseases (Chowdhury *et al.*, 2018).

The source of heavy metal accumulation in the environment is through natural and anthropogenic sources. Anthropogenic activities such as rapid industrialization, overgrowing urbanization and environmental manipulation lead to environmental pollution with heavy metals. Emissions from rapidly expanding industrial areas, mine residues and the accumulation of heavy metals through the disposal of high metal waste pollute the soil and water (Verma *et al.*, 2018).

Arsenic is ubiquitous element, and has many forms on earth. Arsenite and arsenate forms of As are highly toxic to human and animals health. Exposure to As

occurs in many ways. Industrial activities are one of the most important reason of As exposure. Also drinking water and feed may contaminated with As that found in wood preservatives, herbicides, pesticides, fungicides and dyes (Engwa *et al.*, 2019; İriş and Çınar, 2019). Cadmium is another highly toxic metals both for humans and animals. It accumulates in the soil and contaminates pastures (Reis *et al.*, 2010). Cadmium is released into the environment through coal and waste incineration, disposal of metal-containing products, and the use of phosphate fertilizers (İriş and Çınar, 2019). Some plants can accumulate Cd. Therefore, they can be toxic to animals that consume these plants. While the performance of cattle decreases when consuming 5-10 mg/kg of Cd, it affects the health of animals above 30 mg/kg (Reis *et al.*, 2010). Lead is one of the most important contaminants because of its largely use in industrial products. Exposure to lead can occur in many ways such as drinking water, food, cigarettes, industrial processes and domestic sources (Engwa *et al.*, 2019). Because of eating habits cattle reported widely in terms of Pb toxication. Lead toxication mostly affect gastrointestinal, central nervous and hematological systems in living organisms (Thompson, 2007).

Heavy metals can be found as residues in soil, ingestion of plants, sewage wastes, fertilizers and feeds. Apart from these, mine wastes, gasoline with Pb, paints and pesticides are among the causes of metal pollution (Kanbur and Tekeli, 2017). Considering all these reasons, it is important to investigate the level of undesirable substances especially in terms of heavy metals. The aim of this study is to investigate the presence of As, Cd, and Pb metal residues in cattle feed collected from some ruminant farms in Kırıkkale province.

## Materials and Methods

### Sample Collection

The feeds were collected from local enterprises of dairy and beef cattle breeding farms in Kırıkkale Province. A total of 15 total mixed rations samples were collected from the Districts of Kırıkkale Province (4 samples), Yahşihan (3 samples), Karakeçili (1 samples), Balıseyh (1 samples), Delice (2 samples), and Keskin (4 samples). The feed samples were total mixed rations that were prepared by the farm owners. Environmental differences such as the distance of the enterprises from each other were taken into account in the selection of the enterprises. Feed samples were collected in May, 2021. Containers and bags conforming to standards were used for sampling. The samples were stored at +4°C and analyzed in the Nanolab Special Food Control Laboratory for the necessary analysis as soon as possible.

### Sample analysis

The As, Cd, and Pb levels in feed samples were analyzed according to the procedures ISO27085-2009 E modified by Nanolab Special Food Control Laboratory In-house Method- "K.SOP.148" using, inductively coupled plasma mass spectrometry (ICP-MS, AGILENT/7700e, Japan). All chemicals were of analytical reagent quality. Arsenic, Cd, Pb standard stock solutions (1000 mg/L) (Merck) were used (0.05, 0.1, 0.5, 1, 5, 10, 20, 50 µg/L). For calibrations 50, 20, 10, 5, 1, 0.5, 0.1, 0.05 µg/L (ppb), appropriate solutions were made using stock standard solutions.

**Table 1.** Arsenic, cadmium and lead levels in cattle feed samples (mg/kg).

| Metal   | Sample number<br><i>n</i> | Positive<br><i>n (%)</i> | Contamination (mg/kg) |               | Samples exceeding legal level<br><i>n (%)</i> |
|---------|---------------------------|--------------------------|-----------------------|---------------|---|
|         |                           |                          | min-max               | Mean ±SD      |   |
| Arsenic | 15                        | 15 (100%)                | 0.027-0.385           | 0.1475±0.1060 | -   |
| Cadmium | 15                        | 13 (86.67%)              | 0.023-0.053           | 0.0382±0.0079 | -   |
| Lead    | 15                        | 15 (100%)                | 0.050-0.403           | 0.1944±0.1074 | -   |

*n*: sample number, min: minimum level, max: maximum level

Data were given as mean ± SD. The maximum tolerable value is 2 mg/kg for arsenic, 5 mg/kg for lead, 1 mg/kg for cadmium in cattle feed (Republic of Turkey Ministry of Agriculture and Forestry No: 2014/11 Undesirable substances in animal feed)

Approximately 0.5 g of the prepared test sample was weighed into a vessel. Then, 8 ml HNO<sub>3</sub> (%65, Merck) and 2ml H<sub>2</sub>O<sub>2</sub> (%30, Merck) were added, mixed, and burnt at microwave oven for 45 minutes. Samples were cooled and diluted to 50 ml using de-ionized water and shaken several times. It was filtered with cellulose acetate filtrate into a 15 ml falcon tube. After filtrated samples were applied to the ICP-MS device. A blank test tube was also prepared with the same procedure as for feed samples. Arsenic, Cd, and Pb heavy metals in test samples were measured at 75, 111, and 208 mass, respectively. The results were calculated according to the calibration.

The data of the study are given as mean  $\pm$  standard deviation (SD). The evaluation of the data is based on the Republic of Turkey Ministry of Agriculture and Forestry No: 2014/11 Notification of undesirable substances in animal feed.

## Results

Arsenic and Pb residues were found in all of the collected samples, while Cd residues were found in 13 (86.67%) of the collected samples. In the samples with heavy metal residues; the mean value  $\pm$  standart deviation for As was found as  $0.1475 \pm 0.1060$  mg/kg ranging from 0.027-0.385 mg/kg. The mean value  $\pm$  standart deviation for Cd was found as  $0.0382 \pm 0.0079$  mg/kg ranging from 0.023-0.053 mg/kg. The mean value  $\pm$  standart deviation for Pb was found as  $0.1944 \pm 0.1074$  mg/kg ranged from 0.050-0.403 mg/kg. The maximum tolerable limit for As in feed is 2 mg/kg, for Pb in feed stuff is 5 mg/kg, for cadmium in cattle feed is 1 mg/kg (The Republic of Turkey's Ministry of Agriculture and Forestry Announcement No: 2014/11). The results of this study showed that the detected values were below the tolerable limit (Table 1).

## Discussion

Heavy metals are extensively dispersed into the environment along with industrial wastes, as a result, they bio accumulate in the environment. These metals are known as environmental pollutants that pollute natural environments such as water, air and soil, and have toxic effects on humans through animals, plants and the food chain. Arsenic, Cd and Pb are known as the most exposed heavy metals (Yiğit ve Kabakçı, 2018). The main source of As accumulation in agricultural soil is the use of pesticides containing As (Adriano, 2001). Pesticides are commonly used in orchards. High levels of As have been found in pastures that were previously used as orchards (Willett *et al.*, 1993).

Arsenic levels in grain and protein feed mix were reported 10 times higher than those in homegrown feed, probably due to the addition of other ingredients to the feed mix by the feed mill (for example, mineral mixtures containing P) (Li *et al.* 2005). The highest As concentration was reported in mineral mixtures by Li *et*

*al.* (2005). However, acute poisonings with As in cattle are rare (Yiğit and Kabakçı, 2018). In a study conducted in China, the average As content in cattle feed samples was found to be between 0.80-1.38 mg/kg (Zhang *et al.*, 2012). Cang *et al.* (2004) reported 0.13 mg/kg average As content in cattle feed in Jiangsu province. Wang *et al.* (2013) determined an average value of 0.3 mg/kg As in 35 dairy cow feeds in Jiangsu province of China. Nicholson *et al.* (1999) reported an average of 0.37 mg/kg in pelleted dairy cattle feed, 0.49 mg/kg in pelleted beef cattle feed, <0.10 mg/kg As in oat-barley mixture collected from the Wales province of England. Li *et al.* (2005) determined As values of 433  $\mu$ g/kg in dairy feed, 490  $\mu$ g/kg in dairy feed (substitute) and 450  $\mu$ g/kg in heifer feed. In the presented study, it was revealed that As levels ( $0.1475 \pm 0.1060$  mg/kg) were higher than cattle feeds in the Izmir region declared an average value of  $0.046 \pm 0.127$  mg/kg (Güvercin, 2010), and lower than dairy feeds in Jiangsu city of China with an average value of 0.3 mg/kg As (Wang *et al.*, 2013). Arsenic levels of this study did not exceed the legal level recorded as 2 mg/kg.

High concentrations of Cd (up to 10 mg/kg) have been found in forages grown in fields near industrial zinc coating areas where urban sludge is used as fertilizer (Smith, 1986). Cadmium is a human carcinogen associated with lung and prostate cancer therefore it should not be found in feed and feedstuff (Hadjey and Trombetta, 2004). According to the legal restrictions in Turkey Cd in cattle feed should not be more than 1 mg/kg. Nicholson *et al.* (1999) found an average of 0.37 mg/kg Cd in dairy cattle feed, 0.49 mg/kg in pelleted cattle feed, and <0.10 mg/kg in oat barley mixture. Li *et al.* (2005) determined an average of 51  $\mu$ g/kg in dairy feed, 159  $\mu$ g/kg in dairy feed (substitute), and 63  $\mu$ g/kg Cd in heifer feed. Li *et al.* (2010) totally collected 210 feed samples from poultry, cattle, pig, and sheep feed and declared that over % 88.6 of the samples contain Cd residue. The authors reported that the average value of Cd residue in 71 cattle feed collected from Beijing and Fuxin cities in China was 2.79 mg/kg. Güvercin (2010) collected 216 cattle feed from 3 different areas (Kemalpaşa, Torbalı and Kiraz) in Izmir province and detected an average of  $0.07 \pm 0.06$  mg/kg Cd. Cerit *et al.* (2007) stated that the mean Cd value was  $2.016 \pm 1.46$  mg/kg in 4 cattle feed samples collected from 4 different regions (Bursa-Istanbul Highway, Bursa-Izmir Highway, Uludag University Faculty of Veterinary Medicine, Demirtaş) in Bursa. Wang *et al.* (2013) determined that there is 0.42 mg/kg Cd in 35 dairy cow feeds in Jiangsu province of China. The occurrence of Cd in feed may be due to the addition of Zn sulfate, phosphates and Zn oxide supplements to the feed as Cd generally found in these mineral supplements (Li *et al.*, 2010). Thus some resaerchers demonstrated 1–3.6% of Cd in the Zn sulfate additive which shows that phosphate and zinc sulfate were the main sources of Cd in the animal compound feeds (Nong, 2002; Zhong and Jiang, 2005). In the presented study, it was revealed that Cd levels ( $0.0382$

$\pm 0.0079$ ) were lower than cattle feeds in Izmir region that declared an average value of  $0.07 \pm 0.06$  mg/kg (Güvercin, 2010) and in Beijing and Fuxin city of China with an average value of 2.79 mg/kg Cd (Li *et al.*, 2010).

Animal species have different susceptibility to Pb poisoning. Cattle and sheep are considered to be more susceptible to Pb poisoning (Şanlı, 2002). Wang *et al.* (2013) determined an average value of 5.2 mg/kg Pb in 35 dairy cow feeds in Jiangsu province of China. Nicholson *et al.* (1999) found an average value of 2.00 mg/kg Pb in pelleted dairy feed, <1.00 mg/kg Pb in pelleted cattle feed, 1.16 mg/kg Pb in the oat-barley mixture, Cerit *et al.* (2007) detected an average value of  $0.902 \pm 0.517$  mg/kg Pb residue in cattle feed. Güvercin (2010) detected an average value of  $0.28 \pm 0.52$  mg/kg Pb, which was higher than the average Pb value of  $0.1944 \pm 0.1074$  mg/kg that was found in this study.

## Conclusion

The results of this study showed that the detected values of As, Cd and Pb were below the tolerable limit for cattle feed. For this reason, it has been revealed that heavy metal pollution in terms of As, Cd and Pb do not pose a risk to use cattle feed. Apart from this, it should be taken into account that metal pollution can be encountered in many situations. A more comprehensive analysis should be made not only of feed but also of other nutrients such as water consumed by the animal. When heavy metals are collectible materials, analyzes in terms of metal pollution in the animal's hair, milk, as well as in blood will contribute to the protection of human and animal health.

## Acknowledgements

We would like to thank NANOLAB Food Control laboratory for supporting this study.

## References

Adriano, D.C. (2001). Trace elements in terrestrial environments: biogeochemistry, bioavailability and risks of metals. Springer, New York.

Balali-Mood, M., Naseri, K., Tahergorabi, Z., Khazdair, M. R., & Sadeghi, M. (2021). Toxic mechanisms of five heavy metals: Mercury, lead, chromium, cadmium, and arsenic. *Frontiers in Pharmacology*, 12: 227.

Baldwin, D.R., & Marshall, W.J. (1999). Heavy metal poisoning and its laboratory investigation. *Ann Clin Biochem*, 36: 267-300

Cang, L., Wang, Y.J., Zhou, D.M., Dong, Y.H. (2004). Heavy metals pollution in poultry and livestock feeds and manures under intensive farming in Jiangsu Province, China. *Journal of Environmental Science*, 16: 371–374

Cerit, H., Mor, F., Kurşun, Ö., & Avanus, K. (2007). Cadmium and Lead in livestock feed and cattle manure from

four agricultural areas of Bursa, Turkey. *Fresenius Environmental Bulletin*, 16 (9b): 1227-1231.

Chowdhury, R., Ramond, A., O'Keefe, L.M., Shahzad, S., Kunutsor, S.K., Muka, T., Gregson, J., Willeit, P., Warnakula, S., Khan, H., Chowdhury, S., Gobin, R., Franco, O.H. & Di Angelantonio, E. (2018). Environmental toxic metal contaminants and risk of cardiovascular disease: systematic review and meta-analysis. *BMJ*, 29;362:k3310. doi: 10.1136/bmj.k3310.

Engwa, G.A., Ferdinand, P.U., Nwalo, F.N., Marian, N., & Unachukwu, M.N. (2019). Mechanism and health effects of heavy metal toxicity in humans. *Intech Open*, DOI: <http://dx.doi.org/10.5772/intechopen.82511>

Garcia, J.D.D. & Arceo, E. (2018). Renal damage associated with heavy metals: review work. *Revista Colombiana de Nefrología*, 5(1): 45-53.

Güvercin, İ. (2010). İzmir ilinde sütlerde bazı ağır metal (Kurşun, Kadmiyum, Arsenik, Cıva, Bakır, Çinko) düzeylerinin belirlenmesi. Ege Üniversitesi Fen Bilimleri Enstitüsü. Doktora tezi, İzmir.

Hardej, D., & Trombetta, L.D. (2004). Metals .In: *Clinical Toxicology Principles and Mechanisms*. Frank A. Barile, CRC PRESS, USA, pp.295-317.

Hejna, M., Gottardo, D., Baldi, A., Dell'Orto, V., Cheli, F., Zaninelli, M., & Rossi, L. (2018). Review: Nutritional ecology of heavy metals. *Animal*, 12(10): 2156-2170.

ISO 27085 (2009). Animal feeding stuffs — Determination of calcium, sodium, phosphorus, magnesium, potassium, iron, zinc, copper, manganese, cobalt, molybdenum, arsenic, lead and cadmium by ICP-AES. International Standart. First edition 2009-04-15.

İriş, C., & Çınar M. (2019). Antioksidan vitaminler ve ağır metal toksisitesi üzerine etkileri. *Bulletin of Veterinary Pharmacology and Toxicology Association*, 10 (3): 135-151

Kanbur, M., & Tekeli, M.Y. (2017). Yem ve yem hammaddelerinde pestisidler, çevre ve besin kirleticileri. *Türkiye Klinikleri Veterinary Science Pharmacology and Toxicology -Special Topics*, 3(2):120-30.

Li, Y., McCrory, D. F., Powell, J. M., Saam, H. & Jackson-Smith, D. (2005) A survey of selected heavy metal concentrations in wisconsin dairy feeds. *Journal of Dairy Science*, 88(8): 2911–2922.

Li, Y., Xiong, X., Lin, C., & Zhang, F. (2010). Cadmium in animal production and its potential hazard on Beijing and Fuxin farmlands. *Journal of Hazardous Materials*, 177 (1-3): 475–480.

Nicholson, F.A., Chambers B.J., Williams, J.R. & Unwin R.J. (1999). Heavy metal contents of livestock feeds and animal manures in England and Wales. *Bioresource Technology*, 70: 23-31.

- Nong, Y.X.(2002). The investigation of cadmium concentration in feed grade zinc sulfate, *Guangxi Agricultural Science* 5: 260 (in Chinese).
- Reis, L.S., Pardo, P.E., Camargos, A.S., & Oba, E. (2010). Mineral element and heavy metal poisoning in animals. *African Journal of Medicine and Medical Sciences*, 1(12): 560-579
- Smith, M. (1986). Effects of long-term, low-level oral cadmium on performance, blood parameters, and tissue and milk mineral concentrations of dairy cattle through first gestation and subsequent lactation, in: Ph.D. Thesis, Pennsylvania State University.
- Şanlı, (2002). Metaller ve inorganik zehirler. In: Veteriner Klinik Toksikoloji. Medipres, Ankara p.: 185-266.
- Thompson, L.J. (2007). Lead. In: *Veterinary Toxicology, Basic and Clinical Principles*. Ed. Gupta RC, Elsevier Inc, USA. pp. 438-442.
- Verma, K., Vijayalakshmy, R., & Chaudhry, V. (2018). Detrimental impacts of heavy metals on animal reproduction: A review. *Journal of Entomology and Zoology Studies*, 6(6): 27-30
- Wang, H., Dong, Y., Yang, Y., Toor, G.S., & Zhang, X. (2013). Changes in heavy metal contents in animal feeds and manures in an intensive animal production region of China. *Journal of Environmental Sciences*, 25(12): 2435–2442.
- Willett, L.B., O'Donnell, A.F., Durst, H.I. & Kurz, M.M. (1993). Mechanisms of movement of organochlorine pesticides from soils to cows via forages. *Journal of Dairy Science*, 76: 1635-1644.
- Yiğit, A. & Kabakçı R. (2018). Çevre kirleticilerden ağır metallerin hayvanlarda hematopoetik sistem üzerine etkileri. *Türkiye Klinikleri Veterinary Science Pharmacology and Toxicology-Special Topics*, 4(1): 9-15.
- Zhang, F., Li, Y., Yang, M., & Li, W. (2012). Content of Heavy Metals in Animal Feeds and Manures from Farms of Different Scales in Northeast China. *International Journal of Environmental Research Public Health*, 9: 2658-2668.
- Zhong, N., & Jiang, J.L. (2005). Advance of studies on the toxicology of cadmium in feedstuff and environment. *Feed Industry* 26: 18–22 (in Chinese).